







Questionnaire -Clinical Challenges 1/2

Please answer the following questions to help us understand your struggles in providing care to patients at risk for or experiencing delirium

1. How many delirious patients do you care for in an average month?

a) none

c) 6-10

b) 1-5

d) more than 10

2. How confident are you identifying delirium?

a) not at all

c) very

b) somewhat

d) completely

c) moderately

3. Which of the following behaviors of patients with delirium do you find challenging? (circle all that apply)

- a) restlessness
- b) disrupting medical apparatus
- c) getting up from bed or chair unassisted
- d) crying out
- e) hallucinations
- f) delusions
- g) disoriented to time, place and circumstance

- h) refusal of care
- i) lethargy
- j) indifference
- k) difficulty participating in medical and personal care activities (eating, bathing, grooming, mobilizing)

4. Which of the following statements describe your challenges surrounding delirium care (circle all that apply)

- a) I really don't know what to do when patients display certain challenging behaviors
- b) I know what to do to address the challenging behaviors, but I don't have enough time to do it
- c) I know what to do to address the challenging behaviors, but I do not have the resources (equipment/items) to do it
- d) I feel comfortable with my knowledge and technique of caring for patients with delirium who have challenging behaviors









Questionnaire -Clinical Challenges 2/2

5. How does your current process of caring for patients at risk for or experiencing delirium affect you? (circle all that apply)

- a) I feel very good about the care I am able to give to these patients
- b) Caring for these patients is the same as caring for any other patient
- c) I feel stressed because these patients may suffer physical or emotional consequences if things don't go right
- d) I feel frustrated to not be able to help these patients avoid or get through this challenging time

6. How do you feel about the family of patients at risk of or experiencing delirium? (circle all that apply)

- a) I would prefer the family not be around as they often get in the way and ask lots of questions
- b) I try to engage the family to participate in care as much as possible
- c) I rely on the family to share information about the patient's cognitive baseline and coping mechanisms
- d) Our team notifies the family if a new delirium occur

Please add any other comments regarding care of patients with delirium:	